





Trilogy™ Pretreatment Solution

TrilogyTM, a Cell MarqueTM patented pretreatment formula, is the ultimate solution for your lab. Unlike other EDTA solutions, TrilogyTM is a near neutral pH solution, making it less harmful on tissue, while still exhibiting the strong unmasking abilities of a conventional EDTA solution. This revolutionary formula combines deparaffinization, rehydration, and unmasking, all in one simple step! TrilogyTM allows for standardization of the pretreatment procedure, which in turn leads to more consistent and reliable results.

50 ml, 20x	920P-04
200 ml, 20x	920P-06
1 liter, ready-to-use	920P-09
1 gallon, ready-to-use	920P-10

Declere™ Pretreatment Solution

Declere™ may be used whenever immunohistochemistry staining is performed on formalin-fixed, paraffin-embedded tissue sections. Using this product encourages standardization of the pretreatment procedure, thereby producing more consistent, more reliable results.

50 ml, 20x92	1P-04
200 ml, 20x	1P-06
1 liter, 20x92	1P-09

Electric Pressure Cooker

Heat induced epitope retrieval (HIER) techniques using $Trilogy^m$ in conjunction with a pressure cooker allows for simultaneous deparaffinization, rehydration, and epitope retrieval.

Staining Dish and Slide Racks

Designed for histological or cytological staining. Each rack vertically holds up to 24 slides. This kit includes two staining dishes and two slide racks.

Diamond: Antibody Diluent

Diamond: Antibody Diluent is an appropriately Tris buffered solution (TBS) for the dilution of both polyclonal and monoclonal antibodies as well as for the preparation of negative control reagents in IHC. It is designed to minimize the non-specific reaction that may be caused by the antibody and encourage specific antigen-antibody binding.

50 ml, ready-to-use	8B-03
200 ml, ready-to-use	8B-05
1 liter, ready-to-use93	8B-09

Emerald: Antibody Diluent

Emerald: Antibody Diluent is an appropriately phosphate buffered solution (PBS) for the dilution of both polyclonal and monoclonal antibodies as well as for the preparation of negative control reagents in IHC. It is designed to minimize the non-specific reaction that may be caused by the antibody and encourage specific antigen-antibody binding. Emerald: Antibody Diluent can also be used to stabilize diluted antibodies when stored at 2-8°C.

50 ml, ready-to-use936	B-03
250 ml, ready-to-use	B-08
1 liter, ready-to-use	B-09